

# Protocol

## Treatment of Status Asthmaticus with Epinephrine

Status asthmaticus is a severe asthma attack that does not respond to the usual therapies. Although reversible, it is a life-threatening issue, because it can lead to respiratory failure and cardiac arrest. Status asthmaticus requires immediate treatment.

Status asthmaticus is caused by severe bronchospasm, in which the smooth muscles of the bronchi suddenly contract and narrow the airways. The symptoms include extreme difficulty with breathing, which causes restlessness and anxiety. Although coughing and wheezing are common symptoms of asthma, a child with status asthmaticus may not cough or wheeze because there is not enough airflow. Advanced symptoms include little or no breath sounds, inability to speak, bluish tinge to the skin (cyanosis), and heavy sweating. If left untreated, status asthmaticus can lead to unconsciousness and cardiopulmonary arrest.

It is within the scope of practice for professional registered nurses in the state of North Carolina to be able to recognize the symptoms of status asthmaticus and to administer epinephrine to persons who suffer the symptoms of status asthmaticus.

### **Protocol:**

All school nurses will have epinephrine available for use in the schools.

### **Management Guidelines for Treating Status Asthmaticus in the schools:**


1. In the event of an asthma episode by student, staff member or visitor to the school, the school nurse will:
  - A. Obtain a history of asthma
  - B. Move patient to a quiet area if possible.
  - C. Administer fast acting inhaler, such as Albuterol, if available. Stock albuterol inhalers may also be used if available in the school (see protocol for stock inhaler use).
  - D. Offer cool water and observe patient for improvement or worsening condition.
  - E. Contact or have another staff member contact parent/guardian, or emergency contact to inform of above.
  
2. If an asthma episode progresses to the point of status asthmaticus, the school nurse will:

- A. Administer epinephrine according to the following approved procedure.
- B. Call 911 or have another staff member do so for immediate transport to an emergency room.
- C. Inform parent/guardian/emergency contact of emergent condition.

Prescribed Dosages for Epinephrine:

- a) Epinephrine 0.3 mg via auto injector for individuals 66 pounds or greater.
- b) Epinephrine (Jr) 0.15 mg via auto injector for individuals 33-66 pounds.

07/06/2022  
Date

  
\_\_\_\_\_  
Nancy Bizzell, MD  
Medical Consultant  
Union County Public Schools